
Tuesday 19 02 2019


Published 18 02 2019, 17:00



AM




PM











Increase in danger of gliding avalanches and wet snow slides as a consequence of warming during the day and solar radiation. Weakly bonded old snow requires caution.

A latent danger of gliding avalanches exists. This applies on steep grassy slopes. As a consequence of warming during the day and the solar radiation, the likelihood of gliding avalanches being released will increase in particular on steep sunny slopes below approximately 2600 m. Medium-sized gliding avalanches are possible. In addition there is a danger of wet loose snow avalanches. This applies in the afternoon, especially on extremely steep southeast, south and southwest facing slopes below approximately 2600 m. Weak layers near the ground can still be released in isolated cases especially on very steep shady slopes, this applies in particular in case of a large load. Weak layers in the old snowpack can be released in isolated cases and mostly by large additional loads also on very steep sunny slopes, in particular in the afternoon.

Snowpack

Danger patterns

(dp 2: gliding snow)

dp 1: deep persistent weak layer

Outgoing longwave radiation during the night will be good. The surface of the snowpack has frozen to form a strong crust and will soften during the day. This applies at low altitude as well as on very steep sunny slopes in particular below approximately 2600 m. Isolated avalanche prone weak layers exist in the old snowpack.


Tendency

Slight decrease in danger of gliding avalanches and wet snow slides as the temperature drops.



Published 18 02 2019, 17:00





Slight increase in avalanche danger as a consequence of warming during the day.

A clear night will be followed in the early morning by quite favourable conditions generally. As a consequence of warming during the day and solar radiation there will be an increase in the danger of wet and gliding avalanches. Avalanches can in isolated cases be released by small loads and reach medium size. The avalanche prone locations are to be found at transitions from a shallow to a deep snowpack above the tree line. This applies in particular on steep shady slopes and adjacent to ridgelines and in gullies and bowls. Backcountry tours should be started and concluded early.

Snowpack

Danger patterns


dp 10: springtime scenario

The surface of the snowpack will freeze to form a strong crust and will soften during the day. Faceted weak layers exist in the bottom section of the snowpack in particular in shady places that are protected from the wind. Only a little snow is lying.

Tendency

A generally favourable avalanche situation will prevail.





Increase in danger of gliding avalanches and wet snow slides as a consequence of warming during the day and solar radiation. Weakly bonded old snow requires caution.

A latent danger of gliding avalanches exists. As a consequence of warming during the day and the solar radiation, the likelihood of gliding avalanches and moist snow slides being released will increase in particular on steep sunny slopes below approximately 2600 m. Weak layers near the ground can still be released in isolated cases especially on very steep shady slopes, this applies in particular in case of a large load.

Snowpack

Danger patterns

(dp 2: gliding snow)

dp 1: deep persistent weak layer

Outgoing longwave radiation during the night will be good. The surface of the snowpack has frozen to form a strong crust and will soften during the day. This applies at low altitude as well as on very steep sunny slopes in particular below approximately 2600 m. Isolated avalanche prone weak layers exist in the old snowpack.

Tendency

Slight decrease in danger of gliding avalanches and wet snow slides as the temperature drops.