








### **Danger Level 2 - Moderate**



# Regular avalanche bulletins with hazard maps will appear again from around the start of December, depending on the snow situation.

The Avalanche Warning Service currently has only a small amount of information that has been collected in the field.

The wind slabs of the last three days have bonded quite well with the old snowpack in particular on southeast to south to southwest facing aspects. The avalanche prone locations are to be found in particular adjacent to ridgelines and in gullies and bowls in all aspects. Mostly avalanches are medium-sized and can be released also by a single winter sport participant. The number and size of avalanche prone locations will increase with altitude.

The snowpack will become increasingly moist. In the regions exposed to heavier precipitation and on steep sunny slopes small to medium-sized wet snow slides and avalanches are to be expected as a consequence of warming during the day and solar radiation.

#### Snowpack

**Danger patterns** 

dp 10: springtime scenario


Over a wide area 10 to 40 cm of snow, and even more in some localities, has fallen in the last three days above approximately 2000 m. Individual weak layers exist in the old snowpack on steep shady slopes, especially above approximately 2800 m. At low altitude no snow is lying.

## Tendency

Increase in danger of moist and wet avalanches as a consequence of warming during the day and solar radiation. High altitudes and the high Alpine regions: Wind slabs are to be evaluated critically.



## **Danger Level 2 - Moderate**



# Regular avalanche bulletins with hazard maps will appear again from around the start of December, depending on the snow situation.

The Avalanche Warning Service currently has only a small amount of information that has been collected in the field. As a consequence of warming during the day and solar radiation more small and, in isolated cases, medium-sized moist and wet avalanches are possible. More frequent moist loose snow avalanches are to be expected. This applies above approximately 2000 m on extremely steep slopes. The prevalence of avalanche prone locations will increase with altitude. Caution is to be exercised in particular in the regions exposed to heavier precipitation.

High Alpine regions: In addition the fresh wind slabs especially adjacent to ridgelines are prone to triggering in some locations. In regions exposed to the foehn wind avalanche prone locations are more prevalent and exist in all aspects.

#### Snowpack

**Danger patterns** dp 10: springtime scenario dp 6: cold, loose snow and wind


Some snow will fall in some regions. The wind will be moderate to strong. In very isolated cases wind slabs are lying on soft layers. This applies in particular on shady slopes above approximately 3000 m. Outgoing longwave radiation during the night will be reduced in some places. The snowpack will become increasingly moist. At low altitude no snow is lying. At intermediate altitudes hardly any snow is lying. The old snowpack will be in most cases stable.

## Tendency

As a consequence of warming during the day and solar radiation there will be an increase in the danger of moist and wet avalanches.



#### **Danger Level 1 - Low**



Regular avalanche bulletins with hazard maps will appear again from around the start of December, depending on the snow situation.

The Avalanche Warning Service currently has only a small amount of information that has been collected in the field.

The snowpack remains in most cases moist. Moist and wet avalanches are possible even now.

#### Snowpack

**Danger patterns** 

dp 10: springtime scenario

The snowpack will be wet all the way through over a wide area. Outgoing longwave radiation during the night will be reduced. At low altitude no snow is lying.

## **Tendency**

Slight increase in danger of moist and wet avalanches as a consequence of warming during the day and solar radiation.