


| 1   | 2        | 3            | 4    | 5         |
|-----|----------|--------------|------|-----------|
| low | moderate | considerable | high | very high |







# Weakly bonded old snow requires caution. Fresh wind slabs adjacent to ridgelines.

Weak layers in the old snowpack can be released in isolated cases by individual winter sport participants. Such avalanche prone locations are to be found in particular on very steep shady slopes above approximately 2000 m, especially at the base of rock walls and behind abrupt changes in the terrain, as well as on wind-loaded slopes. Here medium-sized avalanches are possible.

In addition the mostly small wind slabs adjacent to ridgelines and in pass areas are prone to triggering. Individual avalanche prone locations are to be found in particular on north and east facing slopes at elevated altitudes. The fresh wind slabs are to be avoided in particular in terrain where there is a danger of falling.

## Snowpack

Danger patterns

(dp.1: deep persistent weak layer)

(dp.6: cold, loose snow and wind)


Avalanche prone weak layers exist in the old snowpack, especially on steep shady slopes above approximately 2000 m. Fresh wind slabs are lying on soft layers in particular on near-ridge shady slopes. Only a small amount of snow is lying for the time of year.

# Tendency

Monday: Significant warming. Increase in danger of moist and wet snow slides.







## Weakly bonded old snow represents the main danger.

Weak layers in the old snowpack can be released in some places by individual winter sport participants. The avalanche prone locations are to be found in particular on steep west, north and east facing slopes above approximately 2000 m, especially in shady places that are protected from the wind. Avalanches can reach medium size. Isolated whumpfing sounds can indicate the danger. Meticulous route selection is recommended.

In addition the mostly small wind slabs are prone to triggering in some locations. They are sometimes covered with new snow and are therefore difficult to recognise. Caution is to be exercised adjacent to ridgelines and in pass areas on very steep north and east facing slopes at high altitudes and in high Alpine regions. At elevated altitudes the avalanche prone locations are more prevalent.

#### Snowpack

Danger patterns

(dp.1: deep persistent weak layer )

 $\left( \, \mathsf{dp.6:\,cold,\,loose\,snow\,and\,wind} \, 
ight)$ 

Avalanche prone weak layers exist in the old snowpack, especially on steep shady slopes above approximately 2000 m, as well as on steep sunny slopes in high Alpine regions. Released avalanches and field observations confirm the unfavourable bonding of the snowpack.

As a consequence of new snow and a moderate wind, mostly small wind slabs formed especially adjacent to ridgelines and in pass areas. These are lying on soft layers in particular on shady slopes. Only a small amount of snow is lying for the time of year.


# Tendency

Monday: Significant warming. Increase in danger of moist and wet snow slides. Weak layers in the old snowpack necessitate caution.





#### **Danger Level 2 - Moderate**



## Wind slabs and weakly bonded old snow represent the main danger.

Weak layers in the old snowpack can be released in some places by individual winter sport participants. The avalanche prone locations are to be found in particular on steep west, north and east facing slopes above approximately 2000 m.

The wind slabs of the last few days can be released by a single winter sport participant in some cases in all aspects. Caution is to be exercised in gullies and bowls, and behind abrupt changes in the terrain, as well as adjacent to ridgelines at high altitudes and in high Alpine regions.

Individual medium-sized avalanches are possible.

On very steep sunny slopes individual moist loose snow avalanches are to be expected as the day progresses, but they will be mostly small.

Individual gliding avalanches are possible. This applies especially on steep grassy slopes. Meticulous route selection is advisable.

#### Snowpack

Danger patterns

(dp.1: deep persistent weak layer ) ( dp.6: cold, lc

ver  $\left.
ight
angle
ight.$  ( dp.6: cold, loose snow and wind  $\left.
ight
angle$ 

Avalanche prone weak layers exist in the old snowpack, especially on steep shady slopes above approximately 2000 m, as well as on steep sunny slopes in high Alpine regions. As a consequence of new snow and a sometimes moderate wind from variable directions, further wind slabs formed especially adjacent to ridgelines and in gullies and bowls as well as at high altitude.

These are lying on soft layers in particular on shady slopes. The older wind slabs are lying on surface hoar in some places.

Only a small amount of snow is lying for the time of year.

# Tendency

Monday: Significant warming. Increase in danger of moist and wet snow slides. Wind slabs and weakly bonded old snow represent the main danger.

