
Avalanche.report Wednesday 14.02.2024


Published 13 02 2024, 17:00








ΡM











Old wind slabs require caution. As a consequence of warming and solar radiation, the activity of moist avalanches will appreciably increase.

More recent wind slabs can be released even by a single winter sport participant. In isolated cases the avalanches are medium-sized. The avalanche prone locations are to be found in particular on steep slopes above approximately 2000 m, and adjacent to ridgelines and in gullies and bowls.

As a consequence of warming during the day and solar radiation moist snow slides and avalanches are possible as the day progresses. This applies in particular on steep grassy slopes below approximately 2400 m especially above the tree line. Avalanche prone locations for moist avalanches are to be found on steep southeast, south and west facing slopes. Mostly avalanches are only small.

Snowpack

Danger patterns

dp.6: cold, loose snow and wind

The new snow of the last few days is very loosely bonded in particular on northwest to north to east facing aspects and generally at intermediate and high altitudes. The fresh wind slabs are poorly bonded with the old snowpack in all aspects and at elevated altitudes.

Sunshine and high temperatures will give rise from late morning to softening of the snowpack in particular on sunny slopes below approximately 2400 m.








Tendency

Sunshine and high temperatures will give rise as the day progresses to moistening of the snowpack over a wide area. As a consequence of solar radiation individual moist and wet avalanches are to be expected.





Danger Level 1 - Low



Old wind slabs require caution. As a consequence of warming and solar radiation, the activity of moist avalanches will increase.

More recent wind slabs can be released even by a single winter sport participant. In isolated cases the avalanches are medium-sized. The avalanche prone locations are to be found in particular on steep slopes above approximately 2000 m, and adjacent to ridgelines and in gullies and bowls.

As a consequence of warming during the day and solar radiation moist snow slides and avalanches are possible as the day progresses. This applies in particular on steep grassy slopes below approximately 2400 m especially above the tree line. Avalanche prone locations for moist avalanches are to be found on steep southeast, south and west facing slopes. Mostly avalanches are only small.

Snowpack

The new snow of the last few days is very loosely bonded in particular on northwest to north to east facing aspects and generally at intermediate and high altitudes. The fresh wind slabs are poorly bonded with the old snowpack in all aspects and at elevated altitudes.

Sunshine and high temperatures will give rise from late morning to softening of the snowpack in particular on sunny slopes below approximately 2400 m.

Tendency

The weather conditions will facilitate a gradual stabilisation of the snowpack. Sunshine and high temperatures will give rise as the day progresses to moistening of the snowpack over a wide area.

